

A *Swift* GAZE INTO THE 2006 MARCH 29TH BURST FOREST OF SGR 1900+14G.L. ISRAEL¹, P. ROMANO^{2,3,4}, V. MANGANO⁴, S. DALL'OSSO^{5,1}, G. CHINCARINI^{2,3}, L. STELLA¹, S. CAMPANA², T. BELLONI², G. TAGLIAFERRI², A.J. BLUSTIN⁶, T. SAKAMOTO⁷, K. HURLEY⁸, S. ZANE⁶, A. MORETTI², D. PALMER⁹, C. GUIDORZI^{2,3}, D.N. BURROWS¹⁰, N. GEHRELS⁷, H.A. KRIMM^{7,11}

1. INAF – Osservatorio Astronomico di Roma, Via Frascati 33, I-00040 Monteporzio Catone (Roma), Italy; gianluca, stella@mporzio.astro.it

2. INAF – Osservatorio Astronomico di Brera, Via Bianchi 46, I-23807 Merate (Lc), Italy

3. Università degli Studi di Milano, Bicocca, Piazza delle Scienze 3, I-20126, Milano, Italy

4. INAF – Istituto di Astrofisica Spaziale e Fisica Cosmica Sezione di Palermo, via Ugo La Malfa 153, I-90146 Palermo, Italy

5. Dipartimento di Fisica “Enrico Fermi”, Università di Pisa, Largo B. Pontecorvo 3, I-56127 Pisa, Italy

6. Mullard Space Science Laboratory, University College London, Holmbury St. Mary, Dorking Surrey, RH5 6NT, UK

7. NASA/Goddard Space Flight Center, Greenbelt, MD 20771, USA

8. Space Sciences Laboratory, University of California at Berkeley, 7 Gauss Way, Berkeley, California, 94720-7450, USA

9. Los Alamos National Laboratory, Los Alamos, NM, 87545, USA

10. Department of Astronomy & Astrophysics, Pennsylvania State University, University Park, PA 16802, USA and

11. Universities Space Research Association, 10211 Wincopin Circle, Suite 500, Columbia, MD 21044

Received: 21 December 2007; Accepted: 23 May 2008

ABSTRACT

In 2006 March the Soft Gamma-ray Repeater SGR 1900+14 resumed its bursting activity after about 2 years of quiescence. The *Swift* mission observed the source several times in order to monitor its timing and spectral properties. We report on the intense burst “forest” recorded on 2006 March 29 which lasted for ~ 30 s, when *Swift* was pointing at the source with the narrow field of view instruments. More than 40 bursts were detected both by BAT and by XRT, seven of which are rare intermediate flares (IFs): several times 10^{42} ergs were released. The BAT data were used to carry out time-resolved spectroscopy in the 14–100 keV range down to 8 ms timescales. BAT and XRT simultaneous data (in the 1–100 keV range) were used to characterize the broad-band energy spectra of IFs and verify the results obtained from the BAT-only spectral fits.

This unique dataset allowed us to test the magnetar model predictions such as the magnetically trapped fireball and the twisted magnetosphere over an unprecedented range of fluxes and with large statistics (in terms of both photons and IFs). We confirmed that a two blackbody component fits adequately the time-resolved and integrated spectra of IFs. However, Comptonization models give comparable good reduced χ^2 . Moreover, we found: i) a change of behavior, around $\sim 10^{41}$ erg s⁻¹, above which the softer blackbody shows a sort of saturation while the harder one still grows to a few times 10^{41} erg s⁻¹; ii) a rather sharp correlation between temperature and radii of the blackbodies ($R^2 \propto kT^{-3}$), which holds for the most luminous parts of the flares (approximately for $L_{\text{tot}} \geq 10^{41}$ erg s⁻¹). Within the magnetar model, the majority of these findings can be accounted for in terms of thermalised emission from the E-mode and O-mode photospheres. Interestingly, the maximum observed luminosity coming from a region of ~ 15 km matches the magnetic Eddington luminosity at the same radius, for a surface dipole field of $\sim 8 \times 10^{14}$ G (virtually equal to the one deduced from the spindown of SGR 1900+14).

We also studied the persistent emission of SGR 1900+14 preceding and following the burst “forest” in search for transient and permanent timing and spectral variations on short (days/week) timescales. We found that the timing/spectral property variations are correlated with the flux and are likely only due to changes of the high energy part of the spectrum (above 5 keV), further supporting the twisted “magnetosphere” scenario.

Subject headings: pulsar: individual (SGR 1900+14) — star: flare — star: neutron — X-rays: burst

1. INTRODUCTION

Soft Gamma-ray Repeaters (SGRs) are a small class of high energy transient astrophysical sources characterized by their emission of short (~ 100 ms) bright (10^{39} – 10^{42} erg s⁻¹) bursts of soft γ rays. SGRs have been associated with persistent X-ray counterparts with 1–10 keV luminosity of $\sim 10^{34}$ – 10^{35} erg s⁻¹ and energy spectra that are generally well fit with a power-law model ($\Gamma \sim 2$). A blackbody component with $kT \sim 0.5$ keV has been found in SGR 1900+14 and SGR 1806–20 (Woods et al. 1999; Woods et al. 2004; Mereghetti et al. 2005). In contrast to the common short bursts, three SGRs have emitted one very powerful long-duration burst. These “giant flares” are distinguished by their extreme energies (10^{44} – $10^{46.5}$ erg s⁻¹), their hard spectra at the onset, and the presence of coherent 5–7 s pulsations at the spin period of the neutron star during the decaying tail lasting several minutes.

Both SGR 1806–20 and SGR 1900+14 were found to spin

down secularly at a rate of $\sim 10^{-11}$ to 10^{-10} s s⁻¹ (Kouveliotou et al. 1998, 1999). Combining SGR periods and their first derivatives, and assuming that the $B_{\text{dip}} \sim 3.2 \times 10^{19} (P\dot{P})^{1/2}$ G relationship generally applied to radio pulsars holds for SGRs as well, magnetic field strengths of 10^{14} – 10^{15} G are inferred. The latter values provide strong evidence that SGRs are highly magnetized neutron stars, i.e., “magnetars” (Duncan & Thompson 1992). In the magnetar model, the magnetic field is the dominant source of free energy (orders of magnitudes higher than the rotational energy of the star), powering the persistent emission through low-level seismic activity and heating of the stellar interior (Thompson & Duncan 1996). When magnetic stresses build up sufficiently to crack a patch of the neutron star crust, the resulting “crustquake” ejects hot plasma particles into the magnetosphere, which results in an SGR burst (Thompson & Duncan 1995). Alternatively, the short SGR bursts may arise from magnetic reconnection events in the stellar magnetosphere (Lyutikov 2002).

TABLE 1
SGR 1900+14 SWIFT OBSERVATION LOG FOR POINTINGS CARRIED OUT DURING 2006.

Sequence	Obs/Mode	Start time (UT) (yyyy-mm-dd hh:mm:ss)	End time (UT) (yyyy-mm-dd hh:mm:ss)	Exposure (s)
00202746000 ^a	BAT/EVENT	2006-03-25 20:16:30	2006-03-25 20:17:13	43
00203045000 ^b	BAT/EVENT	2006-03-28 13:50:05	2006-03-28 13:50:48	43
00203109000 ^c	BAT/EVENT	2006-03-29 01:27:53	2006-03-29 01:28:36	43
00203127000 ^d	BAT/EVENT	2006-03-29 02:52:50	2006-03-29 02:53:42	50
00203974000 ^e	BAT/EVENT	2006-04-05 06:43:21	2006-04-05 06:44:04	43
00214277000 ^f	BAT/EVENT	2006-06-10 06:49:02	2006-06-10 07:09:04	1202
00030386001	XRT/PC	2006-03-25 22:50:43	2006-03-27 21:47:47	49564
00030386002	XRT/PC	2006-03-28 01:05:28	2006-03-28 23:48:57	11350
00030386003	XRT/WT	2006-03-29 01:01:00	2006-03-29 23:57:00	18592
00030386004	XRT/WT	2006-03-30 01:07:00	2006-03-30 23:59:58	20055
00030386005	XRT/WT	2006-03-31 01:26:23	2006-03-31 23:59:59	15931
00030386006	XRT/WT	2006-04-01 01:13:58	2006-04-03 13:12:31	45951
00030386008	XRT/WT	2006-04-07 11:48:02	2006-04-07 23:20:00	7846
00030386009	XRT/PC	2006-04-08 00:51:04	2006-04-10 23:38:58	29274
00030386010	XRT/PC	2006-04-11 00:48:12	2006-04-11 23:44:55	18250
00030386011	XRT/PC	2006-04-12 00:47:49	2006-04-12 23:43:57	20503
00030386012	XRT/PC	2006-04-13 00:53:42	2006-04-13 18:41:34	10702
00030386013	XRT/PC	2006-04-14 22:22:41	2006-04-14 23:59:56	1602
00214277000	XRT/PC	2006-06-10 06:54:59	2006-06-10 08:56:03	4071

^aFirst BAT trigger: 2006-03-25 20:16:40.28 UT.

^bSecond BAT trigger: 2006-03-28 13:50:15.00 UT.

^cThird BAT trigger: 2006-03-29 01:28:03.99 UT.

^dFourth BAT trigger: 2006-03-29 02:53:09.46 UT.

^eFifth BAT trigger: 2006-04-05 06:43:31.36 UT.

^fSixth BAT trigger: 2006-06-10 06:53:00.8 UT.

Giant flares likely result from a sudden reconfiguration of the star's magnetic field that produces large fractures in the crust and propagates outwards through Alfvén waves of enormous power. The ~ 0.2 s initial spike of giant flares marks the ejection of the largest part of the total event energy (up to $\sim 10^{47}$ ergs) as well as the onset of a relatively large surface fracture (Schwartz et al. 2005). The detection of deca-, hecto-, and kilo-Hertz QPOs in the decaying phases (tails) of giant flares have been interpreted in terms of global seismic oscillations following the fracture formation (Israel et al. 2005; Strohmayer & Watts 2005). The super-Eddington fluxes observed in the tails of these events are thought to be possible because of the suppression of the electron scattering cross sections in the presence of very strong magnetic fields (Paczynski 1992).

Four confirmed SGRs are known so far, three of which are in our Galaxy and one in the Large Magellanic Cloud. The Galactic SGRs are located very close to the plane of the Galaxy, indicating that SGRs belong to a young stellar population. Furthermore, two of these SGRs are likely associated with clusters of very massive stars (Fuchs et al. 1999; Vrba et al. 2000; Eikenberry et al. 2001).

Burst activity in SGRs and, in particular, in SGR 1900+14 occurs sporadically and is quite diverse in behavior. Burst active phases of SGRs vary both in total energy released and duration, tending to be concentrated into relatively narrow intervals (weeks or months) separated by relatively long periods (years) of quiescence. Since its discovery in 1979, when bursts were detected three times in three days, SGR 1900+14 remained in quiescence until 1992, when a handful of bursts were again detected within a few days (Mazets & Golenetskii 1981; Kouveliotou et al. 1993). In 1998 the source entered an unprecedented level of activity during which more than 1000 bursts were recorded within 9 months, and culminated with the ~ 400 s long and rather intense 1998 August 27 giant flare event ($\sim 4 \times 10^{44}$ erg s $^{-1}$). Burst activity was detected until

1999. Particularly interesting, are two events detected in May and September 1998, during which "bunching" of short and long burst were detected (Hurley et al. 1999; Mazets et al. 1999). Finally, after almost two years of inactivity, on 2001 April 18, an intense and long (~ 40 s) burst was detected from SGR 1900+14 (Guidorzi et al. 2004). The energy released by this flare (few $\times 10^{42}$ ergs) was less than that of the giant flare, though larger than that of the most common short SGR bursts. Consequently the flare was dubbed an "intermediate flare" (Kouveliotou et al. 2001). Indeed, looking back at the burst history of SGR 1900+14, it was realized that a handful of similar events, characterized by a longer duration (few seconds–few tens of seconds) and a higher fluence than those characterizing short bursts, were present. These bursts were then grouped into the class of intermediate flares likely forming a continuum in terms of duration and fluences (Olive et al. 2004).

A number of broad-band spectroscopic studies of short and intermediate flares were carried out in the past by different missions. These found that, above 15–20 keV spectra are usually well fitted by an optically-thin thermal bremsstrahlung (OTTB) with characteristic temperatures ranging from 20 to 40 keV (Aptekar et al. 2001). Laros et al. (1986) and Fenimore et al. (1994) studied the cumulative spectral properties of 100 bursts from SGR 1806–20 in the 5–200 keV energy range finding that the number of the photons in the spectrum of the detected bursts was remarkably stable in spite of the large intensity spread (a factor of 50) and that the low-energy (below 15 keV) data were inconsistent with the back-extrapolation of an OTTB model that provided a good fit to the high-energy portion of the spectrum. Qualitatively similar spectral properties were measured during a bright intermediate flare from SGR 1900+14 using *HETE*–2 (Olive et al. 2004) data, which also showed that the OTTB model largely overestimated the flux at low energies (<15 keV). The broad-band spectrum (7–100 keV) was best fitted by the sum of two




FIG. 1.— 15–100 keV BAT light curves with a time resolution of 1 ms obtained during the burst “forest” of 2006 March 29.

blackbody (BB) laws. Similar conclusions were reported by Strohmayer & Ibrahim (2000) confirming the OTTB limits below 15 keV by using data from the Rossi X-Ray Timing Explorer.

Feroci et al. (2004) analyzed the 1.5–100 keV BeppoSAX spectral properties of 10 short burst from SGR 1900+14, further confirming that the widely used optically-thin thermal-bremsstrahlung model provides acceptable spectral fits for energies higher than 15 keV but severely overestimates the flux at lower energies. A better fit was obtained by means of an alternative spectral model such as two blackbodies or a cut-off power law. Most of these studies were hampered by one or more of the following limitations: relatively poor statistics, narrow energy bandpass, poor time resolution, off-axis data, and low sensitivity of wide field of view instruments.

Cumulative analysis of 50 bursts detected by *HETE-2* from SGR 1900+14 showed the presence of a time lag of about 2.2 ± 0.4 ms between the 30–100 keV and 2–10 keV radiation bands implying a rapid spectral softening and energy re-injection mechanism (Nakagawa et al. 2007). Alternatively, the softer spectral component could be reprocessed radiation from the harder emission which might be generated near the neutron star surface. More generally, the observed time lag favors spectral models with (at least) two components. In any case the temperatures of the 2BB model does not seem to depend on either the burst intensity or morphology.

It has been assessed that burst activity in the SGRs can have a persistent effect on the underlying X-ray source (Woods &

Thompson 2006 and references therein). During the 1998 burst activity phase of SGR 1900+14, the X-ray counterpart became brighter, its energy spectrum was altered, and the pulse shape changed dramatically. A similar behavior was also recorded from SGR 1806–20 when resumed its burst activity phase in 2004 culminating with the 2004 December 27th giant flare, and later during the 2005–2006 decaying phase (Woods et al. 2007). It has also been proposed that the bursting activity phases are related to the magnetosphere. The correlation between the persistent X-ray flux and the spectral hardness observed in several magnetar candidates on timescales of years could be accounted for if the evolution is regulated by varying “twist” of the magnetosphere, as in the case of 1RXS J170849–400910 (Rea et al. 2005a and Israel et al. 2007b) and SGR 1806–20 (Rea et al. 2005b and Woods et al. 2007). As discussed by Thompson et al. (2002), magnetars may differ from standard radio pulsars because their magnetic field is globally twisted inside the star, up to a strength of about 10 times the external dipole and, occasionally, some of its helicity propagates across the NS surface through crustquakes, leading to an impulsive twist up of the external field, as well. Such an evolving magnetic field is responsible for the different forms of activity and also for the glitches. The basic idea of the “twist” scenario is that when a static twist is formed, currents flow into the magnetosphere. As the twist angle $\Delta\phi_{\text{NS}}$ grows, electrons provide an increasing optical depth to resonant cyclotron scattering, leading to the build-up of a flatter photon power-law component.




FIG. 2.— BAT and XRT (WT) light curves obtained simultaneously during the burst “forest” of 2006 March 29. Different energy ranges are shown: 1–4 and 4–10 keV for the XRT (panels X1 and X2, respectively), and 15–25 keV, 25–40 keV, 40–100 keV and >100 keV for the BAT (panels B1, B2, B3, and B4, respectively). The XRT light curves were background-subtracted and corrected for vignetting, PSF losses, as well as for pile-up effects. It is evident from the comparison between XRT and BAT light curves that, on average, the IFs are harder than the short bursts, though notable exceptions are present.




FIG. 3.— Outcoming of the our selection criterion of relying upon a minimum of 4000 photons in each spectrum. For the first three seconds of the 2006 March 29 “forest” we reported the net counts (upper panel) and the integration time (lower panel).

TABLE 2

SUMMARY OF THE AVERAGE REDUCED χ^2 AND STANDARD DEVIATION FOR DIFFERENT SINGLE- AND MULTI-COMPONENT MODELS OBTAINED FOR THE 729 TIME-RESOLVED BAT SPECTRA AND THE 8 BAT+XRT SPECTRA OF THE MOST INTENSE BURSTS.

Spectral Model	$\langle \chi^2_\nu \rangle$ BAT+XRT ^a	σ	$\langle \chi^2_\nu \rangle$ BAT	σ	Parameters (#)
Bremss	4.84	1.17	1.71	1.95	2
DiskBB	2.91	0.84	1.17	0.51	2
CompST	2.41	0.16	1.08	0.42	3
CutoffPL	1.36	0.07	1.07	0.23	3
Bremss+BB	1.33	0.25	1.06	0.28	4
CompTT	0.88	0.07	0.99	0.17	4
BB+BB	0.88	0.08	1.01	0.16	4
Bremss ^b	...	...	0.91	0.11	2

^aWe used a free scaling factor between XRT and BAT of the order of 10%.

^bFit carried out in the 15–50 keV range.

At the same time, returning currents produce an extra heating of the star surface and an increased X-ray flux. The few long-term X-ray monitoring observations of magnetars collected until now are consistent with a scenario in which the twist angle steadily increased before the glitch epochs, culminating with glitches, sometimes bursts, and a period of increased timing noise, and then decreased, leading to a smaller flux and a softer spectrum.

In this paper, we report the results obtained from our analysis of the data collected by the *Swift* Burst Alert Telescope (BAT; Barthelmy et al. 2005) and X-Ray Telescope (XRT; Burrows et al. 2005) which monitored SGR 1900+14 during a burst active phase in 2006 March (Palmer et al. 2006). SGR 1900+14 triggered BAT five times in a few days at the end of 2006 March, and once a couple of months later. In particular, on March 29 (Romano et al. 2006a) an extremely intense emission event (fluence of $\sim 2\text{--}3 \times 10^{42}$ ergs), lasting ~ 30 s, of short and relatively energetic bursts was recorded while both BAT and XRT were simultaneously observing the source thanks to an automatic slew performed in response to a previous trigger. About 40 single bursts were detected during this “forest”, or “storm”, a handful are long enough (≥ 500 ms) to be considered “intermediate flares” (IFs). The unprecedented BAT and XRT statistics allowed us to perform both average and time-resolved spectral analysis in the 1–100 keV band, the first ever on a large sample of intermediate flares on ms-timescales. We also studied the persistent emission of SGR 1900+14 preceding and following the “forest” in search for transient and permanent spectral variations on timescales of days and weeks, timescales which have been studied rarely. The timing properties of the event are discussed in an accompanying paper (D. Palmer et al. 2008, in preparation).

2. OBSERVATIONS AND DATA REDUCTION

Table 1 reports the log of the *Swift* observations of SGR 1900+14 that were used for this work. The BAT data of the bursts were accumulated in event mode, with full temporal and spectral resolution, and archived in sequences labeled by the BAT trigger number (the first six sequences in Table 1). The sequence corresponding to the March 29 “forest” is 00203127000. The XRT data of pointed observations were archived in the following 13 sequences of Table 1, which span from 2006 March 25 to 2006 June 10. Note that the XRT data of the “forest” are stored in sequence 0030386003.

The BAT data were analyzed using the standard BAT analysis software distributed within FTOOLS v6.0.5. The arrival times of events were converted to arrival times in the




FIG. 4.— Time-resolved results from the spectral fits obtained in the case of the $BB+BB$ model for the first three seconds of the BAT light curve. The soft BB temperature and radius, the hard BB temperature and radius are reported together with the observed flux in the 10–100 keV band and the reduced χ^2 (from the top to the bottom panel).

reference frame of the Solar System Barycenter (SSB) using the `barycorr` task. Mask weighted (i.e. background subtracted) light curves were created from event data in the 15–25, 25–40, 40–100, and 15–100 keV energy ranges at 1 ms and 4 ms time resolution (see Figure 1 and the four lower panels of Figure 2 for the light curves of the March 29 “forest”). Response matrices for BAT spectra were generated with the task `batdrngen` using the latest spectral redistribution matrices. For our spectral fitting (`XSPECv12.2.1`) we considered the 14–100 keV energy range and applied an energy-dependent systematic error vector¹. The time resolution of BAT is 100 microseconds, so pile-up is never an issue under normal circumstances (see Barthelmy et al. 2005). No deadtime effects have been found for these observations (BAT team; private communication).

The XRT data were first processed by the Swift Data Center at NASA/GSFC into Level 1 products (calibrated and quality-flagged event lists). Then they were further processed with the `Heasoft` (v6.0.5) to produce the final cleaned event lists. In particular, we ran the task `xrtpipeline` (v0.10.3) applying standard filtering and screening criteria, i.e., we cut out temporal intervals during which the CCD temperature was higher than -47°C , and we removed hot and flickering pixels which are present because the CCD operated at a temperature higher than the design temperature of -100°C due to a failure in the active cooling system. An on-board event threshold of ~ 0.2 keV was also applied to the central pixel, which has been proved to reduce most of the background due to either the bright Earth limb or the CCD dark current (which depends on the CCD temperature).

For our analysis we considered Windowed Timing (WT) and Photon Counting (PC) mode data (see Hill et al. 2004 for

a full description of read-out modes) and further selected XRT event grades 0–12 and 0–2 for WT and PC data, respectively (see Burrows et al. 2005). The arrival times of events were converted to arrival times in the reference frame of the SSB.

During sequence 00030386003 (i.e. the “forest”) the count rate of the outburst was high enough to cause pile-up even in the WT mode data. Therefore, to account for this effect, the WT data were extracted in a rectangular 40×20 -pixel region with a 4×20 - or 6×20 -pixel region excluded from its center depending on the peak source intensity, i.e. with a 4×20 -pixel exclusion region during the first and third snapshot² (where the observed count rate reached ≈ 350 counts s^{-1}), and with a 6×20 -pixel exclusion region during the second snapshot (observed count rate reaching ≈ 1300 counts s^{-1}). The size of the exclusion region was determined following the procedure illustrated in Romano et al. (2006b) and correspond to 30–39% (6–4 pixel hole) of the XRT PSF. Ancillary response files were generated with the task `xrtmkarf` within `FTOOLS`, and account for different extraction regions and PSF corrections. We used the latest spectral redistribution matrices in the Calibration Database maintained by HEASARC.

The PC data show an average count rate of ~ 0.08 counts s^{-1} throughout the entire monitoring campaign, therefore no pile-up correction was necessary. We extracted the source events in a circle with a radius of 20-pixels ($\sim 47''$). To account for the background, we extracted WT events within a rectangular box (40×20 pixels), and PC events within an annular region (radii 85 and 110 pixels) centered on the source and far from background sources.

The energy-resolved light curves during the burst active phase is shown in Figure 2 both for the XRT (panels marked with X1 and X2) and the BAT (panels B1 to B4). The XRT

¹ http://heasarc.gsfc.nasa.gov/docs/swift/analysis/bat_digest.html

² A snapshot is a continuous pointing at the target.




FIG. 5.— Square of radii of the two blackbodies as a function of their temperatures for the whole ~ 30 s BAT dataset. Red filled squares and dark green filled circles mark the BB_s component for luminosities below and above $3 \times 10^{40} \text{ erg s}^{-1}$, respectively. Violet filled triangles and blue filled stars are the same quantities as above for the BB_h component. Black diamonds marks the Olive et al. (2004) measurements obtained for the 2001 IF. We also plot two representative power-laws for $R^2 = kT^\alpha$, with α equal -3 and -4 (the latter corresponding to the expected relation for a pure BB.)

light curves were background-subtracted and corrected for vignetting and PSF losses, as well as for pile-up.

2.1. Time-resolved BAT Spectroscopy

This section refers to the analysis of the BAT dataset recorded after the fourth trigger (sequence 00203127000; see Table 1). In consideration of the extremely pronounced variability of the source during the burst active phase we adopted the following strategy for the time-resolved spectroscopic analysis: we selected a 4000-count threshold for the accumulation of each spectrum. This resulted in a set of 729 BAT mask weighted (i.e. background subtracted) spectra extracted from the event data of sequence 00203127000. The choice of the threshold is our best tradeoff at optimizing the S/N of the spectra, while still closely following the sharpest features of the light curve. Figure 3 shows the results of our spectral accumulation criterion for the first three seconds of the BAT light curve. In particular, the choice of 4000 counts per spectrum is such that (i) the rapid increase of the first IF is well sampled, (ii) the first two short bursts (peaking at about 3.10 s and 3.35 s) are resolved in time, (iii) 90% of the IF decay is sampled with a time resolution of ≤ 10 ms. All spectral channels with poor statistics (below 20 counts) were removed before fitting. The resulting accumulation time of the BAT spectra ranges from 8 ms to about 400 ms.

Several different spectral models were adopted in the fit of

the whole sample of 729 spectra in an automatic fashion. For each spectrum and model we recorded the values of the best-fit parameters, unabsorbed flux, and reduced χ^2 . Then, we averaged the measured χ^2 and inferred their standard deviations. The latter quantities can be considered as a qualitative indication of the spectral model fit goodness, on average. The results are given in Table 2 ordered with the increasing number of degrees of freedom in the adopted model.

Similarly to previous studies reported above and in the literature, we tested both single and multi-component models. Former models include: an OTTB (*Bremss*) given that it was found to well describe the hard X-ray spectra of SGR bursts, a power-law with an exponential cut-off (*CutoffPL*) which may be considered an extension of the simple OTTB, a disk blackbody (*DiskBB*) which is a generalization of a thermal spectrum including also geometrical parameters, and different versions of a Comptonized spectrum (such as *CompST*) representative of models involving a thermal spectrum modified by mechanisms which preserve the photon number. Among double component models we considered a two blackbody (*BB+BB*) able to fit well the broad band energy spectra of both short and long SGR burst, an OTTB with a soft blackbody in order to adjust the OTTB model extrapolation at low energy, and a Comptonization model where soft photons are up-scattered in a hot plasma taking into account relativistic effects (*CompTT*; Titarchuk 1994) which is, as a first approx-

TABLE 3

XRT WT AND BAT SPECTRAL FIT RESULTS OF THE 7 IDENTIFIED IFs AND ONE SHORT BURST (SB) FOR THE $BB+BB$ MODEL. FLUXES ARE NOT CORRECTED FOR THE N_H .

Bin	N_H (10^{22} cm^{-2})	kT (keV)	R_{BBs} (km) ^a	kTh (keV)	R_{BBs} (km) ^a	$F_X^{1-10 \text{ keV}}$ ($\times 10^{-7} \text{ ergs cm}^{-2} \text{ s}^{-1}$)	$F_X^{10-100 \text{ keV}}$ ($\times 10^{-7} \text{ ergs cm}^{-2} \text{ s}^{-1}$)	χ^2_{red}	d.o.f.
IF1	2.0 ± 0.3	6.2 ± 0.3	15.9 ± 4.2	12.0 ± 0.3	3.5 ± 1.2	4.68 ± 0.15	103.1 ± 1.8	0.84	135
IF2	1.9 ± 0.3	5.6 ± 0.3	17.6 ± 4.7	10.4 ± 0.3	3.6 ± 1.5	4.77 ± 0.15	64.4 ± 1.7	1.15	153
SB	1.2 ± 0.3	5.5 ± 0.3	13.4 ± 4.0	12.1 ± 0.4	2.2 ± 0.8	2.71 ± 0.13	29.1 ± 0.9	0.77	105
IF3	2.1 ± 0.6	5.4 ± 0.3	17.1 ± 5.4	10.2 ± 0.4	3.6 ± 1.6	4.29 ± 0.24	74.5 ± 1.9	0.97	100
IF4	2.1 ± 0.5	6.1 ± 0.3	17.4 ± 4.9	11.7 ± 0.3	3.7 ± 1.4	5.49 ± 0.23	127.8 ± 3.0	0.88	115
IF5	2.8 ± 0.4	6.1 ± 0.3	17.1 ± 4.6	11.0 ± 0.3	4.1 ± 1.6	5.19 ± 0.17	89.8 ± 2.4	1.07	146
IF6	2.3 ± 0.4	5.1 ± 0.3	19.0 ± 5.9	9.8 ± 0.3	3.8 ± 1.6	4.70 ± 0.18	53.5 ± 1.9	0.73	117
IF7	2.7 ± 0.5	5.7 ± 0.3	17.1 ± 5.1	11.6 ± 0.3	3.6 ± 1.2	4.71 ± 0.20	85.1 ± 2.8	0.77	116


^a Assuming a distance to the source of 10 kpc.

FIG. 6.— Time-resolved bolometric luminosity of the BB_{soft} vs. that of the BB_{hard} (in units of 10^{40}). The solid line marks the power-law relation $L_{\text{soft}}^{40} = (L_{\text{hard}}^{40})^\alpha$ with $\alpha = 0.70 \pm 0.03$ (1σ c.l.) obtained by fitting points in the $0.01\text{--}3 L_{\text{soft}}^{40}$ range. Filled squares represent the measurements by Feroci et al. (2004) for short bursts.

imation, a generalization of both the $BB+BB$ model (in the case of saturated Comptonization) and of the *CompST* one.

The procedure of increasing new spectral components (therefore increasing the number of free parameters) was stopped when the F-test probability for the inclusion of an additional component became less than 3σ . It is evident from Table 2 that small (close to unity) reduced χ^2 are obtained, for BAT spectra, for one or two component models with free parameters ≥ 3 . Moreover, small reduced χ^2 are obtained, for BAT+XRT joined spectra, only for two component models with 4 free parameters. The inclusion of a third component (5 or 6 free parameters) is not statistically significant.

The *CompTT* and $BB+BB$ are by far those that gave the best reduced χ^2 (average of 0.99 and 1.01). Moreover, as already reported by several authors, while the *OTTB* model fits the spectra well on relatively narrow energy ranges (reduced $\chi^2 \approx 0.9$ in the 15–50 keV range), it fails to produce an acceptable fit once a broader energy interval is considered (reduced $\chi^2 \approx 1.7$ in the 14–100 keV energy range). A better description is obtained by adding a BB component to the *OTTB* spectrum (reduced $\chi^2 \approx 1.07$). This additional BB component turned out to have a characteristic temperature similar to that of the

softer BB component in the 2BB model.

The average temperatures and sizes of the BB components in the $BB+BB$ during the IFs is $\langle kT_{\text{soft}} \rangle = 4.8 \pm 0.3$ keV with $\langle R_{\text{soft}} \rangle = 30 \pm 2$ km, and $\langle kT_{\text{hard}} \rangle = 9.0 \pm 0.3$ keV with $\langle R_{\text{hard}} \rangle = 5.7 \pm 0.5$ km, which are in good agreement with previous results based on an IF (Olive et al. 2004) and a large sample of short bursts (Nakagawa et al. 2007). However, with respect to the latter cases, the present *Swift* data represent the best opportunity (so far) to study the above reported spectral parameters as a function of time with good statistics coupled to fine timing resolution. Figure 4 shows, as an example, the results of the automatic spectral fitting in the case of the 2BB model for the first 3 s of the high time resolution BAT data, while in Figures 5 and 6, we used the inferred values for the entire 30 s duration and tried to characterize the main properties of the adopted model.

2.2. Time-resolved BAT+XRT Spectroscopy

Joint fit of the BAT and simultaneous XRT/WT data of the bursting phase associated with the fourth trigger was performed with a second set of eight BAT and XRT/WT spectra (sequences 00203127000 and 00030386003). The eight time intervals during which these spectra were accumulated correspond to the occurrence of the seven IFs plus a canonical short burst (as a comparison). We report the main spectral parameters in Table 2 and Table 3.

The results of the XRT plus BAT fits (see Table 2) confirm those obtained with the BAT in the previous section: the *CompTT* and the $BB+BB$ models gave the best reduced χ^2 (≈ 0.9), the *OTTB* (Bremss Table 2 and hereafter) the worst. However, when a BB component is added to the latter component a better solution is obtained (with a number of d.o.f. equal to that of the *CompTT* and the $BB+BB$ models, though the reduced χ^2 (≈ 1.3 ; Table 2) remains too high to be acceptable. More generally, the addition of a BB component to the other poorly fitted models, such as the *CompST* and *CutoffPL*, results in a smaller reduced χ^2 , though not yet acceptable and with a larger number of free parameters (5). In all cases the characteristic temperature of the soft thermal component is similar to that of the soft BB in the 2BB model. These findings, together with similar results obtained by the analysis of broad band spectra of SGR bursts from other missions, in slightly different (but overlapping) energy intervals, and by using different procedures/algorithms, make us confident that the *Swift* fitting results are reliable.

We note that adopting a different abundance than solar values for the absorption multiplicative component (Anders & Grevesse 1989; Lodders 2003) does not significantly modify the spectral parameters for the joint XRT plus BAT spectral




FIG. 7.— The 1–10 keV Swift XRT spectra (PC mode only) of the persistent emission of SGR 1900+14 are shown together with the model (BB plus PL) residuals. The corresponding *Swift* data sequences are 00030386001 (black lines), 00030386002 (red lines and filled squares), 00030386009 (green lines), and 00030386010–6014 (blue lines and filled circles).




FIG. 8.— The 1, 2, and 3σ confidence regions obtained for the BB temperature kT and the power-law photon index Γ for each spectrum during persistent emission from SGR 1900+14. Note that the temperature kT is consistent with being constant, while Γ is variable, and harder than in quiescence, at more than 3σ level.

fit.

2.3. XRT Monitoring of the Persistent Emission

The approximately two-month time span covered by the *Swift* monitoring observations was also used to study the timing and spectral properties of the persistent emission. Only the PC data were used for spectral analysis in order to rely upon a better signal to noise ratio. WT data were also accumulated and used for timing analysis, after removing bursts, in order to rely upon a higher time resolution. The corresponding *Swift* data sequences used to carry out the spectral analysis are 00030386001 (≈ 50 ks effective exposure




FIG. 9.— The 1, 2, and 3σ confidence regions obtained for the BB and PL normalizations for each spectrum during persistent emission from SGR 1900+14. Similarly to Figure 8 the BB component appears to be approximately constant, while the PL normalization is variable and higher than in quiescence, at more than 3σ level.

time), 00030386002 (≈ 11 ks), 00030386009 (≈ 29 ks), and 00030386010–6013 (≈ 51 ks). In order to minimize the possible burst contamination in PC mode, spectra were accumulated after having excluded ~ 10 s time-interval just before and after each good time intervals (GTIs) corresponding to automatic switches between PC and WT modes. In fact, these switches are likely due to high count rate events from cosmic rays and/or from genuine bursts from SGR 1900+14. A comparison between the filtered and unfiltered spectra did not result in the detection of significant differences, making us confident that burst contamination, if any, is negligible. As in the previous cases, the data were rebinned with a minimum of 20 counts per energy bin to allow χ^2 fitting within XSPEC (v12.2.1). We further removed those energy channels with poor statistics after background subtraction.

We assumed an absorbed power-law plus blackbody model as inferred with deeper *XMM-Newton* observations (Mereghetti et al. 2006). We left the absorption free to vary but forced to be the same in different spectra and obtained $N_H = (2.3 \pm 0.1) \times 10^{22} \text{ cm}^{-2}$. The results of the fits are reported in Table 4, while fitted spectra are shown in Figure 7 together with their residuals expressed in units of σ . There is marginal evidence of parameter variability as a function of observed flux: the spectrum becomes harder for larger flux levels. In order to properly quantify such a spectral variability and its flux dependence, we carried out a bi-dimensional fit (contour plot) of the main spectral parameters which has also the advantage of better taking into account the possible correlations among them. The results of these additional fits are shown in Figure 8 and 9. The ellipsoidal shape of the 1, 2, and 3σ confidence regions (the innermost region corresponding to 1σ) clearly indicates that parameters (and, more generally, the BB and PL components) correlate with each other. Additionally, the two-parameter fits confirm the spectral evolution (c.l. $> 3\sigma$) of the persistent spectrum of SGR 1900+14 as a function of flux: the source displayed harder spectra just before (March 28) and after (April 8) the burst “forest” event (March 29), and returned back to the pre-event level in mid-April. These results are similar to those observed in other magnetar candidates, such as SGR 1806–20 among SGRs and

1RXS J170849–400910 among AXPs, the main difference being in the timescales on which these spectral variations occurred: years for the latter two sources and days/weeks for SGR 1900+14. In the previous cases this trend has been modelled in terms of “twisting” of the magnetosphere (Thompson et al. 2002).

The WT and PC XRT data were also used to look for the ~ 5.2 s pulsations of SGR 1900+14. Given the limited XRT statistics in the persistent component and the relatively small pulsed fraction of the pulses (~ 15 – 20% level; semi-amplitude of modulation divided by the mean source count rate) we used the period value, $P=5.19987\pm 0.00007$ s, inferred from an *XMM-Newton* observation carried out on 2006 April 1 (Mereghetti et al. 2006). We detected the SGR 1900+14 pulsations on 2006 March 29, the same day as burst “forest” event, in two time intervals preceding (March 29 from 01:02 to 01:26 UT) and following (from 04:22 to 04:42 UT) the “forest” by approximately 1 hr (in both cases the signal was detected at about 3σ c.l.). By merging all the XRT observations carried out in March/April (from 30 March 01:07 to 7 April 23:20) and by using $\dot{P}=(9.2\pm 0.4)\times 10^{-11}$ s s $^{-1}$ reported by Mereghetti et al. (2006), the pulsations could be detected at a high significance level. Figure 10 summarizes the results of the timing analysis. Both the pulse fractions and shapes inferred from the first dataset are different from those obtained from the second and third datasets, and, more generally, from what observed on average since the discovery of this source. At the same time, the pulse shapes of the second and third datasets are significantly different. The value and shape of the the latest dataset is virtually equal (to within the uncertainties) to those obtained by Mereghetti et al. (2006) about 3 days after the BAT event.

2.4. UVOT Observations and Data Reduction

The Swift UV/Optical Telescope (Romig et al. 2005) is a 30 cm modified Ritchey-Chretien telescope co-aligned with the XRT. It features a photon-counting micro-channel plate intensified CCD detector optimized for UV and optical light over the 170–600 nm range; its six broadband colour filters sample this range.

The UVOT was taking data in the *V* filter (central wavelength 5460 Å) during the March 29 outbursts of SGR 1900+14. No optical counterpart is detected in the resulting image, accumulated between 2006-03-29 02:39:57 and 2006-03-29 03:06:00 UT with a total exposure time of 1539 s. In the following we derived an upper limit to the optical emission during the burst “forest”. Counts were extracted from a $2''$ radius aperture at the position of the source in the XRT; the size of the aperture was chosen so as to minimize contamination by a $V \sim 18.6$ mag star (USNO number 0975-14353360) which is only $3''$ from the XRT position. The counts were aperture-corrected up to $6''$, and used to calculate a 3σ *V*-band magnitude upper limit for any counterpart of 20.7 mag. Adopting the Akerlof et al. (2000) estimate for A_V of ~ 10 mag for the Galactic reddening in the direction of SGR 1900+14 we estimate a dereddened upper limit of 10.7 mag.

For each UVOT image coincident with an outburst, counts were extracted from $4''$ radius apertures at the position of the source in the XRT. The counts were aperture-corrected up to $6''$ (optical filters) or $12''$ (UV filters); the different apertures are due to the broader PSF in the UV, and used to calculate 3σ magnitude upper limits for any counterpart. Using the Akerlof et al. (2000) estimate of $A_V \sim 10$ magnitudes for




FIG. 10.— The 1–10 keV *Swift* XRT lightcurves of SGR 1900+14 from approximately one hour before and after the BAT event (2006 March 29; top and mid panels), and the first week of April 2006 (bottom panel). In all cases we used $P=5.19987\pm 0.00007$ s, as determined from an *XMM-Newton* observation carried out on 2006 April 1 (Mereghetti et al., 2006). For the about one week-long baseline of the April 2006 lightcurve dataset we used $\dot{P}=(9.2\pm 0.4)\times 10^{-11}$ s s $^{-1}$. The pulsed fraction, semi-amplitude of modulation divided by the mean source count rate, is reported on the right of each panel.

the Galactic reddening in the direction of SGR 1900+14, we corrected the upper limits for Galactic extinction assuming $E(B-V)=3.125$. The resulting values are listed in Table 5.

3. DISCUSSION

Swift recorded a rather intense and rare series of short bursts and intermediate flares (the densest part of which lasted for ~ 30 s) from SGR 1900+14 during 2006 March exploiting its rapid follow-up pointing capability. This allowed us to carry out, for the first time, a detailed study of the average and time-resolved spectral properties of this source during such events. Moreover, the continuous monitoring of SGR 1900+14 on longer timescales, afforded a study of the changes of the spectral and timing properties of the persistent emission.

3.1. Burst emission

We used several different spectral models comprising one or two components, to fit both the time-resolved BAT spectra and the time-averaged (over a whole single burst or IF) BAT+XRT spectra. Interestingly, only two (*CompTT* and 2BBs) out of the five models considered (*CompST*, *CutoffPL*, *Bremss+BB*, *CompTT* and 2BBs) which fit the BAT spectra well, are also able to model the BAT+XRT spectra (see Table 2 and Table 3). An optically-thin thermal bremsstrahlung (*Bremss*) does not yield an acceptable reduced χ^2 in either cases (BAT and BAT+XRT spectra). This result is consistent with previ-

TABLE 4

XRT PC SPECTRAL FIT RESULTS OF THE PERSISTENT EMISSION OF SGR 1900+14 DURING THE FOUR TIME INTERVALS REPORTED IN THE TEXT AND BY ASSUMING THE $BB+PL$ MODEL. FLUXES ARE NOT CORRECTED FOR THE N_H VALUE (WHICH IS LEFT TO VARY BUT CONSTANT AMONG SPECTRA).

Interval MJD	kT (keV)	R_{BBs} (km) ^a	Γ	$F_X^{1-10\text{keV}}$ ($\times 10^{-12}$ ergs cm ⁻² s ⁻¹)	χ^2_{red}	d.o.f.
53819.9519–53821.9082	0.39 ± 0.02	3.4 ± 0.5	2.4 ± 0.1	4.6 ± 0.8	1.13	125
53822.0455–53822.9923	0.51 ± 0.06	2.9 ± 0.6	1.9 ± 0.3	6.3 ± 1.7	0.91	39
53833.0355–53835.9854	0.48 ± 0.03	3.3 ± 0.4	2.1 ± 0.2	5.0 ± 1.4	0.96	86
53836.0335–53840.0	0.38 ± 0.02	4.3 ± 0.8	2.3 ± 0.1	5.0 ± 0.7	0.94	137

^a Assuming a distance to the source of 10kpc and an absorption hydrogen column of $2.3 \pm 0.1 \times 10^{22}$ cm⁻².

TABLE 5

THE TIMES OF SGR 1900+14 OUTBURSTS OCCURRING WHEN UVOT WAS TAKING DATA, WITH THE FILTER IN USE AT THE TIME (CENTRAL WAVELENGTH IN BRACKETS), EXPOSURE TIME OF THE IMAGE IN SECONDS, 3σ MAGNITUDE UPPER LIMIT AT THE XRT POSITION.

Flare time	Filter	Exp	UL
2006-03-25 23:34:07.748	UVM2 (2200 Å)	642	21.3
2006-03-26 00:59:22.680	V (5460 Å)	213	18.9
2006-03-26 13:30:04.620	B (4350 Å)	216	20.8
2006-03-26 16:50:17.632	UVW2 (1930 Å)	851	20.9
2006-03-28 20:32:31.456	UVM2 (2200 Å)	261	20.8
2006-03-29 02:53:09.46	V (5460 Å)	1539	20.7

ous spectral analyses carried out on broad band spectra of a few short bursts and a 4 s-long IF (see, Feroci et al. 2004 and Olive et al. 2004). Notably, the *Bremss* model describes the BAT spectra well only when a reduced energy interval, 15–50 keV, is considered. In the following we discuss in greater detail the results inferred from the 2BB model (note that the *CompTT* model shape is virtually identical to that of the 2BB in the limit of saturated Comptonization). Moreover, in consideration of the difficulties in explaining short bursts and IFs in the accretion scenario (they have extreme super-Eddington luminosities are in fact observed), we discuss here the implications of our results within the magnetar scenario.

With respect to the former studies of the broad-band spectrum of similar events, the *Swift* observation of 2006 March 29 has the advantage of a number of events (>40 between short bursts and IFs), therefore largely improving the statistics over which the detailed analysis can be carried out. The impact of the increased statistics and time resolution is evident when comparing the *Swift* results with previous ones. On 2001 July 2, the *HETE-2* instruments recorded a 3.5 s IF from SGR 1900+14 and a time resolved ($\Delta t > 30$ ms) spectral analysis was carried out in the 2–100 keV range (Olive et al. 2004). They found that the 2BB model fitted the spectra well, the higher temperature BB evolving in a manner consistent with a shrinking emitting region, the lower temperature BB showing a constant radius. They also suggested that, within the magnetar model, the 2BB model might be an approximation of a more complex multi-temperature spectrum. Their Figures 5 and 7 can be easily compared with our Figures 4 and 5: the *Swift* data fill in those regions on the kT - R^2 plane, which remained unexplored with *HETE-2*.

Several new properties can be immediately inferred: a) the *Swift* data populate, almost homogeneously, all temperatures between ~ 2 and 12 keV, and the *HETE-2* measurements (black diamonds in Figure 5) can be regarded as a subset of them; b) the distribution is such that a sharp edge between the populated regions and the rest of the kT - R^2 plane is clearly present, with a cut-off (or sharp turn) in the distribution of

R_{BBh} at $kT_h \approx 12$ –13 keV. This sharp edge provides an estimate of the typical size of the relevant emitting regions (by assuming a reference distance of 10 kpc): 30–200 km range for the BB_s component and 3–30 km range for the BB_h one. c) there is an additional turn in the BB_h component size between 6 and 16 km, at approximately 10 keV (see Figure 5), and d) there is a strong correlation between kT and R_{BB}^2 corresponding to the brightest phases of the IFs, which is consistent with a power-law with index of about -3 (while, were the luminosity the same for all events, the relationship expected for a BB component would be $y=x^{-4}$, where $x = kT$ and $y = R_{BB}^2$).

The sharp edge in the distribution of the 2BB parameters suggests that a saturation effect occurs. In order to investigate this effect further, we divided the data points of the two BB components in two samples, below and above a bolometric luminosity of $L = 3 \times 10^{40}$ erg s⁻¹ (see Figure 5). The luminosity threshold was chosen so as to roughly separate the peaks of the bursts and flares from the inter-burst/flare time intervals. We note that the *HETE-2* data lie just above and below the flux threshold, but never reach the values corresponding to the peaks and tops of the *Swift* IFs. As we can see from Figure 5, as long as each BB component is below $\sim 3 \times 10^{40}$ erg s⁻¹, their two luminosities are strongly correlated ($L(BB_{hard})$ vs $L(BB_{soft})$ is well fitted by a power law with index 0.7 ± 0.3 at 1σ c.l.). This correlation, which seems indicative of a physical link between the two BB components, was already noticed by Feroci et al. (2004) who analyzed the time-averaged spectral properties of 10 short bursts from SGR 1900+14 recorded by *BeppoSAX* in April 2001 (filled squares in Figure 6). It is apparent that the *BeppoSAX* measurements are fully consistent with those obtained with *Swift* and can be regarded as a subsample of the latter. On the other hand, this trend was not reported by Olive et al. (2004) for the IF observed by *HETE-2* in 2001 (empty diamonds in Figure 6). Intriguingly, the correlation no longer holds above $\sim 3 \times 10^{40}$ erg s⁻¹ for the *Swift* data. Above this value, the luminosity of the BB_s component reaches a maximum in the $(7\text{--}14) \times 10^{40}$ erg s⁻¹ range, while that of the BB_h continues to increase up to $\sim 3 \times 10^{41}$ erg s⁻¹. Clearly, this previously unknown behavior strengthens the idea of a saturation mechanism in the burst emission.

The emission properties of the higher luminosity flares ($L > 3 \times 10^{40}$ erg s⁻¹), combined with the spectral information, places some basic constraints on their production mechanism. The possible location of the emitting region and the main processes likely involved in shaping the spectra have been discussed by Thompson & Duncan (1995, hereon TD95). Lyubarsky (2002) studied the propagation of a trapped fireball through the ultramagnetized magnetosphere, and computed an approximate emission spectrum for it. However, the comparison of these predictions with the results from our observations or those by Olive et al. (2004) is not straightforward.

ward. As already noted by Olive et al. (2004), the “modified blackbody” proposed by Lyubarsky (2002) predicts a significant depression below the blackbody of the higher energy tail, which is not found in the data (the fit remains good up to energies ~ 100 keV). Clearly, more detailed spectral models for the burst are required.

In the following, we will concentrate on new important information (mainly concerning the brightest part of IFs) which has been inferred from our spectral analysis. In particular, we discuss in detail three main findings: i) the existence of two blackbody components, the harder one ($kT_h \sim 7\text{--}11$ keV) having a systematically smaller radius ($R_{BBh} \leq 25$ km) than the softer one ($R_{BBs} \sim 25\text{--}100$ km); ii) the existence of a clear correlation between the luminosities of the two components up to $\sim 3 \times 10^{40}$ erg s $^{-1}$, above which value the luminosity of softer blackbody shows signs of saturation, while that of the harder blackbody still grows up to a few times 10^{41} erg s $^{-1}$; iii) the existence of a correlation between temperature and blackbody radius, which holds for the most luminous parts of the flares (approximately for $L_{\text{tot}} > 4 \times 10^{40}$ erg s $^{-1}$).

The first finding shows that different thermal components originate from different regions around the NS, ranging from its surface (at $R < R_{\text{NS}}$) to well up in the magnetosphere, at a height of several stellar radii. The maximum observed flare luminosity is $\sim 3 \times 10^{41}$ erg s $^{-1}$, attained by the hard blackbody component at an effective temperature of ~ 10 keV and radius of ~ 15 km. Interestingly, this matches well the magnetic Eddington luminosity, $L_{\text{Edd,B}}$, at that same radius, for a surface dipole field $\approx 8 \times 10^{14}$ G (this value of the B-field is very close to that deduced from the spindown rate of SGR 1900+14), where (cfr. Paczynsky 1992, TD95):

$$L_{\text{Edd,B}}(r) \approx 2L_{\text{Edd}} \left(\frac{B}{10^{12}\text{G}} \right)^{4/3} \approx 2 \times 10^{40} \left[\frac{B}{B_{\text{QED}}} \right]^{4/3} \left(\frac{R}{R_{\text{NS}}} \right)^{2/3} \text{ erg s}^{-1}. \quad (1)$$

If radiation originated from a trapped hot fireball this would be in line with the discussion of Thompson & Duncan (1995, 2001), according to which the radiative efficiency of a magnetic confined fireball never exceeds (to within a small factor) the magnetic Eddington flux. In fact, ablation of matter from the NS surface would increase the scattering depth in a super-Eddington radiation field, thus providing a self-regulating mechanism. The radius and temperature that we inferred at maximum burst luminosity are also in good agreement with the prediction for the emission coming from a trapped fireball (TD95, cfr. their discussion for the best fit temperature and radius in the case of SGR 1806–20).

Most remarkably, we find that the luminosity of the soft component can be larger than 10^{40} erg s $^{-1}$ out to $R \sim 100$ km. For $R > 30\text{--}40$ km this value exceeds the magnetic Eddington luminosity at the corresponding radius (for a dipole field geometry) and suggests that magnetospheric confinement plays an active role out to those distances, with magnetic stresses balancing radiation forces. If this were the case, the saturation of the soft component at $\sim 10^{41}$ erg s $^{-1}$ would be related to the magnetic field strength at the relevant radii and its maximum ability to retain the trapped fireball matter which is subject to very high radiation pressure. A detailed investigation of the possible equilibrium configurations is beyond the scope of this paper and will be addressed in a future work.

Our second finding [point ii)] implies that, in the luminosity range $10^{40}\text{--}10^{41}$ erg s $^{-1}$, the total radiation energy is divided almost equally between the two components. Combining the information of Figures 5 and 6 we can see that, for a given

luminosity $\leq 10^{41}$ erg s $^{-1}$, there are two separate emission regions: a smaller and hotter one, whose radiating area suggests emission from (part of) the NS surface with a relatively high effective temperature ($10\text{--}12$ keV), plus a second, possibly magnetospheric region, with significantly larger emitting area and lower effective temperature ($3\text{--}7$ keV). The hotter component attains the highest luminosities, *i.e.* $> 10^{41}$ erg s $^{-1}$, as its radius grows slightly and the effective temperature decreases. On the other hand, the colder component is characterized by a minimum blackbody radius of ~ 25 km and maximum temperature of ~ 7 keV.

A possible interpretation involves the different way in which photons with ordinary (O) or extraordinary (E) polarization mode (a properties introduced by the presence of a birefringent medium, such as the magnetic field; Mészáros et al. 1980) propagate across the magnetosphere. Since the scattering cross section of E-mode photons is much reduced in presence of strong magnetic fields, E-mode photons have a scattering photosphere which is located much closer to the NS than that of O-mode photons. On the other hand, in supercritical magnetic field ($B > B_{\text{QED}}$), E-mode photons have a non-negligible probability of splitting (and switching to the O-mode). The probability is a strong function of energy, so that E-mode photons of high energy cannot travel far from their scattering photosphere before splitting into O-mode photons. On the other hand, O-mode photons will be entrained with streaming electrons and baryons and advected out to where their scattering optical depth becomes $\tau(O) \sim 1$ (TD95; Lyubarsky 2002). As a consequence of Compton scattering (and, with lower efficiency, photon merging), a fraction of O-mode photons can also switch back to the E-mode. Therefore, the two modes are effectively coupled and are advected at comparable rates as long as mode switching is efficient: an approximately equal distribution of energy in the two modes is thus to be expected (TD95; Lyubarsky 2002). This may suggest that the two observed spectral components reflect the population of photons in the two polarization modes and, thus, the regions from which they are emitted.

We note that the minimum radius of the cold blackbody almost corresponds to the maximum radius of the hot component, the distribution of points in the R^2 vs. kT plane being continuous across this point (see Figure 5). This suggests the existence of a narrow zone of separation between the two emission regions. In order to further characterize this zone, we studied the R^2 vs. kT distribution for each of the seven longest IFs and for one short burst we already discussed in previous sections (see also Table 3 and Figure 11). Remarkably, the transition between the two BB components is more evident for the brightest flares (IF1 and IF7 in Figure 11) where the upper left edge of the parameter distribution of the hard component is confined to smaller radii ($R \leq 20$ km) and correspondingly higher temperatures ($kT \geq 8$ keV). The soft component seems less dependent on the flux than the hard one, its minimum size being of the order of $\geq 25\text{--}35$ km in all IFs. Qualitatively, this behavior is in agreement with the above scenario. In fact, a larger brightness can be reached when the burst energy is trapped closer to the NS surface, where the stronger magnetic field allows a larger flux (of E-mode photons) to be released. Moreover, the sharp and narrow gap between the emitting regions of the two BBs, at a radius of ($\sim 20\text{--}30$ km), can be interpreted as the signature of the presence of an E-mode splitting photosphere (TD95); this represents the surface (at radius $R = R_{\text{QED}}$) below which the magnetic field is supercritical ($B \geq B_{\text{QED}}$), photon splitting




FIG. 11.— Same as Figure 5 but for each of the seven longest intermediate flares (IF1 to IF7) plus a short burst (SB) also reported in Table 3.

is efficient and the two polarization modes can remain coupled. E-mode photons can stream freely to the observer from the splitting photosphere (at radius $R_{\text{QED}} \sim 20\text{--}30\text{ km}$), which then would naturally define a maximum size for the emission region of radiation in this polarization mode. On the other hand, O-mode photons could not originate from regions internal to the splitting photosphere, as their scattering optical depth is still much larger than unity within that volume.

If this scenario is correct, it implies that the emission region of E-mode photons can range from the base of the trapped fireball (near the NS surface) to a slightly higher region in the magnetosphere, but well within the splitting photosphere. In this case, part of the internal energy of the coupled E and O-mode photons is lost in the E-mode photosphere (adiabatic) expansion from near the NS to the splitting photosphere. This degrades the original spectrum towards lower temperatures and larger radii, up to R_{QED} . On the other hand, O-mode photons might either be released near the splitting photosphere or advected to much larger radii, depending on the value of their optical depth at that location (see TD95). In any case an improved treatment of the above effects would be required in order to develop a self-consistent model.

Finally, our third finding is the apparent correlation between the surface and temperature of the spectral components at different luminosities. This behavior does not have a clear interpretation. As shown in Figure 5, the top part of the data strip is well fit by a power-law $\Sigma \propto T_{\text{BB}}^{-3}$, where Σ is the radiating blackbody surface. Since $\Sigma T_{\text{BB}}^4 = L$, this implies $T_{\text{BB}} \propto L$. Therefore, along lines of constant luminosity, the radiating surface would scale like T^{-4} , i.e. steeper than the data points (on the edge of the allowed region in the kT vs R^2 plane). The $\Sigma T^3 = \text{const}$ relation that implies a roughly constant number of emitted photons per unit time.

At the highest luminosities ($\geq 10^{41} \text{ erg s}^{-1}$) of the flares, where the softer blackbody saturates, the observed correlation involves both components. It is tempting to attribute this to the fact that, in this luminosity range, Comptonization, and/or adiabatic losses play a dominant role in the formation of the emerging spectrum, both processes being characterized by conservation of the photon number (photon splitting should have only a minor impact). This is in agreement with the prediction by TD95, that at effective temperatures less than $\sim 10\text{ keV}$ photon splitting would not be efficient in maintaining a pure Planck spectrum for both modes. However, if the photon chemical potential is sufficiently small compared to the photon temperature, deviations from a Planck spectrum are expected only at low energies (TD95; Lyubarski 2002), which we do not observe given they are masked by absorption.

On the other hand, at luminosities $\leq 10^{41} \text{ erg s}^{-1}$, the two blackbodies approximately share the same luminosity, therefore the colder one emits a larger number of photons. This suggests that efficient photon number-changing processes (such as photon splitting or double Compton scattering) may play a key role in the formation of the spectrum. Indeed, we note that the photon number ratio is roughly the same as the temperature ratio in this case, since the luminosities are approximately equal. As this ratio is peaked around 2 and extending from 1.5 and 4, it appears to be qualitatively compatible with the combined effects of photon splitting and double Compton scattering (cfr. Lyubarski 2002).

Finally, we briefly comment on the sharp turn in the hard component distribution at around $kT \sim 10\text{ keV}$ and subsequent

cut-off at $\sim 12\text{--}14\text{ keV}$ which, together with the $\Sigma T^3 = \text{const}$ relation, are among the most distinctive properties of Figure 5. Among others, these features clearly imply that no thermal component with characteristic temperature above the cut-off is detected, though the BAT energy interval extend up to at least 100 keV . Within the discussed scenario it is tempting to account for the above properties in terms of photon splitting process and, in particular, its efficiency as a function of (mainly) flux and energy. In fact, the turn at $kT \sim 10\text{ keV}$ marks the drift of the parameter distribution from the $\Sigma T^3 = \text{const}$ relation, identifying the passage between photon number-conservation (such as Comptonization) and photon number-changing (photon splitting) emitting processes. As reported by TD95, the minimum temperature at which photon splitting is efficient in maintaining a nearly Planckian spectrum for E-mode photons, is $\sim 10\text{ keV}$.

3.2. The persistent emission

The *Swift* XRT monitoring data allowed us to study the evolution of the persistent emission of the SGR 1900+14 on relatively short timescales starting from one day before up to one an half month after the BAT event. We clearly detected a spectral evolution in the higher energy end of the $1\text{--}10\text{ keV}$ spectrum as a function of flux, the PL component becomes flatter for larger fluxes. The changes in the persistent component are temporary and last for approximately a couple of weeks, while we detected variability on daily timescales. It is also interesting to note that the spectral results we obtained on 2006 April 8 (see P3 interval in Table 4 and the green lines in Figures 7, 8 and 9) are consistent, to within the uncertainties, with those reported by Mereghetti et al. (2006) from a deeper *ToO XMM-Newton* observation carried out on 2006 March 1.

Both AXPs and SGRs have shown a correlation between the X-ray flux and the spectral hardness (see, for example, Woods et al 2007; Rea et al. 2005a); this has been explained in terms of the onset of a “twist” in the magnetosphere. As discussed by Thompson et al. 2002, magnetars may differ from standard radio pulsar because their magnetic field is globally twisted inside the star, up to a strength of about 10 times the external dipole and, occasionally, this field is expected to fracture the crust and lead to a twist of the external field. The basic idea is that when such a twist is formed, currents flow into the magnetosphere. As the twist angle $\Delta\phi_{\text{NS}}$ grows, electrons provide an increasing optical depth to resonant cyclotron scattering, leading to the build up of a flatter photon power-law component. At the same time, larger returning currents produce an extra heating of the star surface and increased X-ray flux. Therefore, both increased activity and glitches are expected to be associated with erratic fracturing of the crust. Observations obtained until 2003 were consistent with a scenario in which the twist angle was steadily increasing before the glitch epochs, culminating in glitch(es) and period of increased timing noise, and then decreasing, leading to a smaller flux and a softer spectrum. The decay time of the global twist is given by (Thompson, Lyutikov & Kulkarni 2002)

$$t_{\text{decay}} \simeq 40 \Delta\phi_{\text{N-S}}^2 \left(\frac{L_X}{10^{35} \text{ erg s}^{-1}} \right)^{-1} \left(\frac{B_{\text{pole}}}{10^{14} \text{ G}} \right)^2 \left(\frac{R_{\text{NS}}}{10 \text{ km}} \right)^3 \text{ yr}, \quad (2)$$

where $\Delta\phi_{\text{N-S}}$ is the net twist between the north and south magnetic hemispheres, B_{pole} is the magnetic field strength at the pole(s). By using the inferred persistent unabsorbed luminosity of SGR 1900+14 ($\sim 10^{35} \text{ erg s}^{-1}$), the observed time interval, of the order of <1 month, over which the spectral param-

eters change and return to the pre-burst “forest” event, and assuming a standard size for the neutron star surface and a $10^{14} \text{ G} < B_{\text{pole}} < 10^{15} \text{ G}$ we infer a twist angle in the few 10^{-3} to few 10^{-2} radians interval, which is similar to that inferred for the bursting AXP 1E 2259+586 ($\sim 10^{-2}$ rad; Woods et al. 2004).

We regard as unlikely that the hardening and softening of the power-law component (in the 1–10 keV band) is due to a change of the flux and/or photon index of the additional power-law component detected by *INTEGRAL* in the 20–200 keV band flux of several AXPs and SGRs (Kuiper et al. 2006; the SGR 1900+14 emission above 20 keV was re-discovered also by PDS on board *BeppoSAX*; Götz et al. 2006; Esposito et al. 2007). This is implied by the finding that the spectrum of SGR 1900+14 can be accounted for by only one power-law component covering both the soft and hard X-rays (from ~ 1 to ~ 200 keV).

We monitored the pulse shape and pulse fraction of the ~ 5.2 s spin period modulation through a timing analysis of the persistent component. Large changes were detected as a function of flux (see Figure 10): pulses appear to be less pulsed but more structured (to within the uncertainties) and with smaller relative amplitude for increasing fluxes. Such a flux dependence of the pulse properties was already detected in other SGRs and AXPs and related to changes in the persistent emission component. Among SGRs, SGR 1806–20 showed a decrease in the pulsed fraction together with the appearance of additional peaks in the pulse profile after the giant flare of the 2004 December 27 (Rea et al. 2005b, Woods et al. 2007). More recently, a similar behavior was observed in the transient AXP CXOU J164710.2–455216 in correspondence to the onset of an outburst (Israel et al. 2007a). Given the almost constant properties of the BB component, variations are likely due to the PL component which is supposed to originate into the magnetosphere. In particular, The pulse shape changes may indicate an evolution of the geometry of the magnetosphere. Alternatively, the magnetosphere may randomize the outcoming pulsed photons, temporarily decreasing the pulsed fraction.

4. CONCLUSIONS

On 2006 March 29, after several days of burst activity, the Soft γ -ray Repeater SGR 1900+14 displayed a series of short bursts and intermediate flares lasting for about 30 s, during which all the *Swift* instruments were pointed to the source. The good statistics, the fine time resolution (4 ms) and the wide energy coverage (0.2–100 keV) allowed us to carry out a detailed study of the timing and spectral properties during both the bursts and persistent emission. Large variations were detected and their evolution studied over a range of timescales. Although not all the observational properties can be accounted for by the current magnetar scenario, the large majority of them are in overall good agreement with it. In particular, we found:

- a break, around $\sim 10^{41} \text{ erg s}^{-1}$, in the known correlation between the luminosities of the two blackbody components which fit well the BAT time-resolved and BAT+XRT integrated spectra. Above this value the softer blackbody shows signs of saturation, while the luminosity of the harder blackbody still increases up to $\sim 3 \times 10^{41} \text{ erg s}^{-1}$;
- the existence of a correlation between temperature and emitting surface of the blackbodies ($R^2 \propto kT^{-3}$), which

holds through the most luminous phases of the flares ($L_{\text{tot}} \geq 10^{41} \text{ erg s}^{-1}$).

The above two findings together can be interpreted in terms of the two populations of photons from the O- and E-mode polarizations modes which are foreseen in the magnetar scenario and the regions from which they escape: larger and colder the O-mode photon region, harder and smaller the E-mode one.

Moreover, we note that the maximum observed luminosity from flares ($\sim 3 \times 10^{41} \text{ erg s}^{-1}$) is attained by the harder blackbody component at an effective temperature of ~ 10 keV and radius of ~ 15 km, which is similar to the inferred magnetic Eddington luminosity at the same radius, for a surface dipole field $\simeq 8 \times 10^{14} \text{ G}$ (as deduced from the spindown of SGR 1900+14).

Finally, based on our analysis, we do not see any significant difference between IFs and short bursts. Both IFs and short bursts seem to form a continuum in terms of spectral properties (besides duration and fluence).

As far as the persistent emission properties are concerned, we found:

- a clear correlation between the PL photon index Γ and the persistent flux, such that flatter PL components correspond to larger fluxes. The source returned to the usual spectral shape after approximately one week from the burst “forest” event;
- changes in the modulation of the 5.2 s spin period, with the pulse profiles becoming more structured (to within the uncertainties) for increasing persistent flux. The pulse properties returned to their usual state within one week, as well.

The above results are not dissimilar from those observed in other magnetar candidates, although this is the first time that variations are detected on timescales of an hour (for the pulses) and days (for the spectra). These findings are qualitatively in agreement with the expected properties of a “twisted” magnetosphere.

More generally, the unprecedented high quality of the *Swift* data allowed us to gather new important insights into the emission mechanism properties, at a detail level which had not been attained before. We hope that findings like the ones presented in this paper will stimulate a more detailed theoretical treatment of the physics of SGR and AXP bursts.

This work is partially supported at OAR through Agenzia Spaziale Italiana (ASI), Ministero dell’Istruzione, Università e Ricerca Scientifica e Tecnologica (MIUR – COFIN), and Istituto Nazionale di Astrofisica (INAF) grants. This work is supported at OAB and OAR by ASI grants I/011/07/0, I/088/06/0, and at OAB by MIUR PRIN 2005025417. We gratefully acknowledge the contributions of dozens of members of the BAT, XRT, and UVOT Teams at OAB, PSU, UL, GSFC, ASDC, and MSSL and their subcontractors, who helped make the *Swift* mission possible. GLI would like to thank Alaa Ibrahim for useful discussions on the comparison between the burst spectral properties of SGR 1806–20 and SGR 1900+14, Marco Feroci for comparison with burst properties of SGR 1900+14 as observed by *BeppoSAX*, Maxim Lyutikov for comments on the discussion section, and Peter Woods for careful reading of the manuscript and valuable suggestions for its improvement.

Facilities: Swift (XRT). Facilities: Swift (BAT). Facilities: Swift (UVOT).

REFERENCES

- Anders, E. & Grevesse, N., 1989, *Geochimica et Cosmochimica Acta*, 53, 197
- Akerlof, C., et al. 2000, *A&A*, 542, 251
- Aptekar, R. L., Frederiks, D. D., Golenetskii, S. V., Il'inskii, V. N., Mazets, E. P., Pal'shin, V. D., Butterworth, P. S., & Cline, T. L. 2001, *ApJS*, 137, 227
- Barthelmy, S. D., et al. 2005, *Space Science Review*, 120, 143
- Burrows, D. N., et al. 2005, *Space Sci. Rev.*, 120, 165
- Duncan, R. C., & Thompson, C. 1992, *ApJ*, 392, L9
- Eikenberry, S. S., Garske, M. A., Hu, D., Jackson, M. A., Patel, S. G., Barry, D. J., Colonna, M. R., & Houck, J. R. 2001, *ApJ*, 563, L133
- Esposito, P., Mereghetti, S., Tiengo, A., Sidoli, L., Feroci, M., & Woods, P. 2007, *A&A*, 461, 605
- Fenimore, E. E., Laros, J. G., & Ulmer, A. 1994, *ApJ*, 432, 742
- Feroci, M., Caliendo, G. A., Massaro, E., Mereghetti, S., & Woods, P. M. 2004, *ApJ*, 612, 408
- Fuchs, Y., Mirabel, F., Chaty, S., Claret, A., Cesarsky, C. J., & Cesarsky, D. A. 1999, *A&A*, 350, 891
- Götz, D., Mereghetti, S., Tiengo, A., & Esposito, P. 2006, *A&A*, 449, L31
- Guidorzi, C., Frontera, F., Montanari, E., Feroci, M., Amati, L., Costa, E., & Orlandini, M. 2004, *A&A*, 416, 297
- Hill, J. E., et al. 2004, *Proc. SPIE*, 5165, 217
- Hurley, K., Kouveliotou, C., Woods, P., Cline, T., Butterworth, P., Mazets, E., Golenetskii, S., & Frederiks, D. 1999, *ApJ*, 510, L107
- Israel, G. L., et al. 2005, *ApJ*, 628, L53
- Israel, G. L., Campana, S., Dall'Osso, S., Muno, M. P., Cummings, J., Perna, R., & Stella, L. 2007a, *ApJ*, 664, 448
- Israel, G. L., Götz, D., Zane, S., Dall'Osso, S., Rea, N., & Stella, L. 2007b, *A&A*, 476, L9
- Kouveliotou, C., et al. 1993, *Nature*, 362, 728
- Kouveliotou, C., et al. 1998, *Nature*, 393, 235
- Kouveliotou, C., et al. 1999, *ApJ*, 510, L115
- Kouveliotou, C., et al. 2001, *ApJ*, 558, L47
- Kuiper, L., Hermesen, W., den Hartog, P. R., & Collmar, W. 2006, *ApJ*, 645, 556
- Laros, J. G., Fenimore, E. E., Fikani, M. M., Klebesadel, R. W., & Barat, C. 1986, *Nature*, 322, 152
- Lodders, K. 2003, *ApJ*, 591, 1220
- Lyubarsky, Y. E. 2002, *MNRAS*, 332, 199
- Mazets, E. P., et al. 1981, *Ap&SS*, 80, 3
- Mazets, E. P., Cline, T. L., Aptekar', R. L., Butterworth, P. S., Frederiks, D. D., Golenetskii, S. V., Il'inskii, V. N., & Pal'Shin, V. D. 1999, *Astronomy Letters*, 25, 628
- Mereghetti, S., et al. 2005, *ApJ*, 628, 938
- Mereghetti, S., et al. 2006, *ApJ*, 653, 1423
- Mészáros, P., Nagel, W., & Ventura, J. 1980, *ApJ*, 238, 1066
- Nakagawa, Y. E., et al. 2007, *PASJ*, 59, 653
- Olive, J.-F., et al. 2004, *ApJ*, 616, 1148
- Paczyński, B. 1992, *Acta Astronomica*, 42, 145
- Palmer, D. M., Sakamoto, T., Barthelmy, S., Cummings, J., Gehrels, N., Krimm, H., Markwardt, C., & Tueller, J. 2006, *The Astronomer's Telegram*, 789, 1
- Rea, N., Oosterbroek, T., Zane, S., Turolla, R., Méndez, M., Israel, G. L., Stella, L., & Haberl, F. 2005a, *MNRAS*, 361, 710
- Rea, N., Tiengo, A., Mereghetti, S., Israel, G. L., Zane, S., Turolla, R., & Stella, L. 2005b, *ApJ*, 627, L133
- Romano, P., Barthelmy, S. D., Kennea, J. A., Markwardt, C. B., Moretti, A., & Palmer, D. M. 2006a, *GRB Coordinates Network*, 4934, 1
- Romano, P., et al. 2006b, *A&A*, 456, 917
- Roming, P. W. A., et al. 2005, *Space Sci. Rev.*, 120, 95
- Schwartz, S. J., et al. 2005, *ApJ*, 627, L129
- Strohmayer, T. E., & Ibrahim, A. I. 2000, *ApJ*, 537, L111
- Strohmayer, T. E., & Watts, A. L. 2005, *ApJ*, 632, L111
- Thompson, C., & Duncan, R. C. 1995, *MNRAS*, 275, 255 (TD95)
- Thompson, C., & Duncan, R. C. 1996, *ApJ*, 473, 322
- Thompson, C., & Duncan, R. C. 2001, *ApJ*, 561, 980
- Thompson, C., Lyutikov, M. & Kulkarni, S.R. 2002, *ApJ*, 574, 332
- Titarchuk, L. 1994, *ApJ*, 434, 570
- Vrba, F. J., Henden, A. A., Luginbuhl, C. B., Guetter, H. H., Hartmann, D. H., & Klose, S. 2000, *ApJ*, 533, L17
- Woods, P. M., Kouveliotou, C., van Paradijs, J., Finger, M. H., & Thompson, C. 1999, *ApJ*, 518, L103
- Woods, P. M., et al. 2004, *ApJ*, 605, 378
- Woods, P. M., & Thompson, C. 2006, *Compact stellar X-ray sources*, 547
- Woods, P. M., Kouveliotou, C., Finger, M. H., Göğüş, E., Wilson, C. A., Patel, S. K., Hurley, K., & Swank, J. H. 2007, *ApJ*, 654, 470